

Pre-conception, pregnant women and breastfeeding

Each situation should be evaluated on a case-by-case basis, a comprehensive risk assessment should be performed, and women should be encouraged to make as informed a decision as possible regarding YF vaccination

The safety of YF vaccination at the time of conception, during pregnancy and when breastfeeding, has not been systematically evaluated. Each situation should be evaluated on a case-by-case basis, a comprehensive risk assessment should be performed, and women should be encouraged to make as informed a decision as possible regarding YF vaccination in these circumstances.

Pregnancy during travel carries important risks that should be considered carefully prior to booking the trip. More information can be found on our [pregnancy factsheet](#).

Women travelling to YF risk areas and planning a pregnancy should, ideally, wait until 28 days after YF vaccination before conceiving [1]. This time allows for any post-vaccine viraemia (vaccine virus in the mother's blood), which may in theory be a risk to a developing fetus, to settle.

It is important to note that [malaria](#) and [Zika virus](#) transmission often co-exist in YF risk areas. Women who are pregnant should consider postponing their trip to areas where malaria is known to occur [2]. They should also postpone non-essential travel to high risk Zika areas until after pregnancy and should *consider* postponing non-essential travel to moderate risk Zika areas until after pregnancy [3].

The safety of YF vaccine in pregnancy has not been systematically evaluated; a prospective study in Brazil, where women were inadvertently vaccinated in early pregnancy (mean gestational age of 6 weeks) during a YF outbreak, found no increase in fetal malformations, complications to the central nervous system, premature delivery or perinatal deaths [4].

A recent report from the World Health Organization concluded that *'for live vaccines, while there is a theoretical risk to the fetus, no significant adverse outcomes following vaccination have been reported'* [5]. Current UK guidelines advise that pregnant women should be advised not to travel to a high-risk area. When travel is unavoidable, the risk from the disease and the

theoretical risk from the vaccine have to be assessed on an individual basis. Pregnant women should be counselled on the potential benefits and risks of vaccination so that they may make an informed decision [6]. World Health Organization state that in areas where yellow fever is endemic, or during outbreaks, the benefits of vaccination are likely to far outweigh the risk of potential transmission of vaccine virus to the foetus [7].

Inadvertent administration of a vaccine (live or inactivated) during pregnancy does not constitute grounds to terminate the pregnancy [6].

In order to expand the safety data available on vaccines for pregnant women, health professionals are advised to report details of vaccine use during pregnancy to the [UK Teratology Information Service](#) (a group commissioned by UK Health Security Agency). These groups collect pregnancy outcome information from women who have been exposed to drugs and vaccines in pregnancy.

There is conflicting information on whether the immune response to the YF vaccine during pregnancy is impaired [1]. Until further information is available on this, revaccination following pregnancy is recommended for those at continued risk [6]. The timing of the vaccine must also take in to consideration the YF vaccine and breastfeeding information.

Breastfeeding women

Generally, most mothers will wish to, and should be encouraged to continue breastfeeding during travel. Breastfeeding has benefits for both mother and infant; in particular infants who are exclusively breastfed are unlikely to be exposed to other food or drink, which may be contaminated [8].

However, one case of vaccine associated neurologic disease (YEL-AND) in an infant, aged 23 days old [9] and two probable cases (infants five weeks and 10 days old) have been reported in breastfeeding infants since 2009 [10, 11]. All the infants developed YF encephalitis shortly after their mothers received YF vaccine for the first time and during their babies' first month of life. Although all babies were breastfed, the mode of transmission has not been established [11].

Breastfeeding has historically been, and remains, a precaution for YF vaccination and should be avoided where possible. However, if the risk of YF during travel is considered sufficiently high, YF vaccination for the woman who is breastfeeding can be considered.

If a breastfeeding infant is too young to receive YF vaccine and travel cannot be avoided, the parents should be reminded that [insect bite precautions](#) should be scrupulous (e.g. the use of insect repellent appropriate to the infant's age, room screening and mosquito nets). Insect bite precautions should be adhered to, day and night.

Resources

- [Pregnancy factsheet, TravelHealthPro](#)
- [Zika virus, TravelHealthPro](#)
- [Breastfeeding and bottle feeding, TravelHealthPro](#)

REFERENCES

1. Monath, T.P., Gershman M., Staples J.E. Barrett, A.D.T. Yellow fever vaccine, Ch 38 pg 959 in **Vaccines**, 6th Edition edited by Plotkin SA, Orenstein WA Offit, PA 2013, Elsevier Saunders
2. [UK Health Security Agency. Advisory Committee on Malaria Prevention \(ACMP\). guidelines for malaria prevention in travellers from the United Kingdom; last updated 18 January 2023 \[Accessed 20 January 2023\]](#)
3. [TravelHealthPro. Zika: Risk Assessment 27 February 2019 \[Accessed 13 January 2023\]](#)
4. Suzano CES, Amaral E, Sato HK et al. The effects of yellow fever immunization (17DD) inadvertently used in early pregnancy during a mass campaign in Brazil. **Vaccine** 2006; 24(9): 1421-1426
5. [World Health Organization. Meeting of the Strategic Advisory Group of Experts on Immunization. November 2013 – conclusions and recommendations. Wkly Epidemiol Rec 3 Jan 2014 No. 1, 2014, 89, 1–19 \[Accessed 13 January 2023\]](#)
6. [UK Health Security Agency. Yellow Fever. Ch 35. In: Immunisation against infectious disease. January 2020 \[Accessed 13 January 2023\]](#)
7. [World Health Organization. Vaccines and vaccination against yellow fever: WHO position paper. June 2013. Wkly Epidemiol Rec 27, 88; 269–284 \[Accessed 13 January 2023\]](#)
8. [Anstey EH and Shealy K. Travel and breastfeeding. Ch 7 In. Centers for Disease Control and Prevention Yellow book: Health Information for International Travel. \[Accessed 13 January 2023\]](#)
9. [Centers for Disease Control and Prevention. Transmission of yellow fever vaccine virus through breast feeding – Brazil, 2009. MMWR 2010;59\(05\): 130-132. \[Accessed 13 January 2023\]](#)
10. [Kuhn S, Twele-Montecinos L, MacDonald J et al. Case report: probable transmission of vaccine strain of yellow fever virus to an infant via breast milk. 2011 CMAJ; 183, 4. \[Accessed 13 January 2023\]](#)



11. [World Health Organization. Global Advisory Committee on Vaccine Safety. Wkly Epidemiol Rec. 16–17 June 2010. 23 July 2010. 85, 30: 285-292. \[Accessed 13 January 2023\]](#)